

Levels of Undecidability in Infinitary Rewriting: Normalization and Reachability

Jörg Endrullis

Free University Amsterdam, The Netherlands
joerg@few.vu.nl

Abstract. In [2] it has been shown that infinitary strong normalization (SN^∞) is Π_1^1 -complete. Surprisingly, it turns out that infinitary weak normalization (WN^∞) is a harder problem, being Π_2^1 -complete, and thereby strictly higher in the analytical hierarchy.

We assume familiarity with infinitary term rewriting; we further reading we refer to [5,4].

1 Infinitary Strong Normalization and Reachability

Definition 1. A *Turing machine* M is a quadruple $\langle Q, \Gamma, q_0, \delta \rangle$ consisting of:

- finite set of states Q ,
- an initial state $q_0 \in Q$,
- a finite alphabet Γ containing a designated symbol $\square$, called *blank*, and
- a partial *transition function* $\delta : Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$.

A *configuration* of a Turing machine is a pair $\langle q, \text{tape} \rangle$ consisting of a state $q \in Q$ and the tape content $\text{tape} : \mathbb{Z} \rightarrow \Gamma$ such that the carrier $\{n \in \mathbb{Z} \mid \text{tape}(n) \neq \square\}$ is finite. The set of all configurations is denoted Conf_M . We define the relation $\rightarrow_M$ on the set of configurations Conf_M as follows: $\langle q, \text{tape} \rangle \rightarrow_M \langle q', \text{tape}' \rangle$ whenever:

- $\delta(q, \text{tape}(0)) = \langle q', f, L \rangle$, $\text{tape}'(1) = f$ and $\forall n \neq 0. \text{tape}'(n+1) = \text{tape}(n)$, or
- $\delta(q, \text{tape}(0)) = \langle q', f, R \rangle$, $\text{tape}'(-1) = f$ and $\forall n \neq 0. \text{tape}'(n-1) = \text{tape}(n)$.

Without loss of generality we assume that $Q \cap \Gamma = \emptyset$, that is, the set of states and the alphabet are disjoint. This enables us to denote configurations as $\langle w_1, q, w_2 \rangle$, denoted $w_1^{-1}qw_2$ for short, with $w_1, w_2 \in \Gamma^*$ and $q \in Q$, which is shorthand for $\langle q, \text{tape} \rangle$ where $\text{tape}(n) = w_2(n+1)$ for $0 \leq n < |w_2|$, and $\text{tape}(-n) = w_1(n)$ for $1 \leq n \leq |w_1|$ and $\text{tape}(n) = \square$ for all other positions $n \in \mathbb{Z}$.

The Turing machines we consider are deterministic. As a consequence, final states are unique (if they exist), which justifies the following definition.

Definition 2. Let M be a Turing machine and $\langle q, \text{tape} \rangle \in \text{Conf}_M$. We denote by $\text{final}_M(\langle q, \text{tape} \rangle)$ the $\rightarrow_M$ -normal form of $\langle q, \text{tape} \rangle$ if it exists and undefined, otherwise. Whenever $\text{final}_M(\langle q, \text{tape} \rangle)$ exists then we say that M *halts on* $\langle q, \text{tape} \rangle$ with final configuration $\text{final}_M(\langle q, \text{tape} \rangle)$. Furthermore we say M *halts on tape* as shorthand for M *halts on* $\langle q_0, \text{tape} \rangle$.

Turing machines can compute n -ary functions $f : \mathbb{N}^n \rightarrow \mathbb{N}$ or relations $S \subseteq \mathbb{N}^*$. We need only unary functions f_M and binary $>_M \subseteq \mathbb{N} \times \mathbb{N}$ relations.

Definition 3. Let $M = \langle Q, \Gamma, q_0, \delta \rangle$ be a Turing machine with $S, 0 \in \Gamma$. We define a partial function $f_M : \mathbb{N} \rightarrow \mathbb{N}$ for all $n \in \mathbb{N}$ by:

$$f_M(n) = \begin{cases} m & \text{if } final_M(q_0 S^n 0) = \dots q S^m 0 \dots \\ \text{undefined} & \text{otherwise} \end{cases}$$

and for M total (i.e. M halts on all tapes) we define the binary relation $>_M \subseteq \mathbb{N} \times \mathbb{N}$ by:

$$n >_M m \iff final_M(0 S^n q_0 S^m 0) = \dots q 0 \dots$$

Note that, the set $\{>_M \mid M \text{ a Turing machine that halts on all tapes}\}$ is the set of recursive binary relations on $\mathbb{N}$.

We use the translation of Turing machines M to TRSs R_M from [3].

Definition 4. For every Turing machine $M = \langle Q, \Gamma, q_0, \delta \rangle$ we define a TRS R_M as follows. The signature is $\Sigma = Q \cup \Gamma \cup \{\triangleright\}$ where the symbols $q \in Q$ have arity 2, the symbols $f \in \Gamma$ have arity 1 and $\triangleright$ is a constant symbol, which represents an infinite number of blank symbols. The rewrite rules of R_M are:

$$\begin{aligned} q(x, f(y)) &\rightarrow q'(f'(x), y) && \text{for every } \delta(q, f) = \langle q', f', R \rangle \\ q(g(x), f(y)) &\rightarrow q'(x, g(f'(y))) && \text{for every } \delta(q, f) = \langle q', f', L \rangle \end{aligned}$$

together with four rules for ‘extending the tape’:

$$\begin{aligned} q(\triangleright, f(y)) &\rightarrow q'(\triangleright, \square(f'(y))) && \text{for every } \delta(q, f) = \langle q', f', L \rangle \\ q(x, \triangleright) &\rightarrow q'(f'(x), \triangleright) && \text{for every } \delta(q, \square) = \langle q', f', R \rangle \\ q(g(x), \triangleright) &\rightarrow q'(x, g(f'(\triangleright))) && \text{for every } \delta(q, \square) = \langle q', f', L \rangle \\ q(\triangleright, \triangleright) &\rightarrow q'(\triangleright, \square(f'(\triangleright))) && \text{for every } \delta(q, \square) = \langle q', f', L \rangle. \end{aligned}$$

In [2] the TRSs R_M has been extended as follow to prove Π_1^1 -completeness of finiteness of dependency pair problems:

Definition 5 ([2]). For every Turing machine $M = \langle Q, \Gamma, q_0, \delta \rangle$ we define the TRS $R_M^\bullet$ as follows. The signature $\Sigma = Q \cup \Gamma \cup \{\triangleright, \bullet, T\}$ where $\bullet$ is a unary symbol, T is a constant symbol, and the rewrite rules of $R_M^\bullet$ are:

$$\ell \rightarrow \bullet(r) \quad \text{for every } \ell \rightarrow r \in R_M$$

and rules for rewriting to T after successful termination:

$$\begin{aligned} q(x, 0(y)) &\rightarrow T && \text{whenever } \delta(q, S) \text{ is undefined} \\ \bullet(T) &\rightarrow T. \end{aligned}$$

Moreover, we define the TRS R_{pickn} to consist of the following rules:

$$\text{pickn} \rightarrow c(\text{pickn}) \quad \text{pickn} \rightarrow \text{ok}(0(\triangleright)) \quad c(\text{ok}(x)) \rightarrow \text{ok}(S(x)).$$

Proposition 6. *Let M be an arbitrary Turing machine. We define the TRS S together to consist of the rules of $R_M^\bullet \uplus R_{\text{pickn}}$ together with:*

$$\text{run}(T, \text{ok}(x), \text{ok}(y)) \rightarrow \text{run}(q_0(x, y), \text{ok}(y), \text{pickn}), \quad (1)$$

and define a term $t := \text{run}(T, \text{pickn}, \text{pickn})$. Then it holds:

$$\text{SN}_S^\infty \iff \text{SN}_S^\infty(t) \iff >_M \text{ is well-founded.}$$

Proof. See [2]. □

Theorem 7. *Uniform infinitary strong normalization, SN_R^∞ , and for single terms, $\text{SN}_R^\infty(s)$ is Π_1^1 -complete.*

Proof. The Π_1^1 -hardness has been shown in [2] using that well-foundedness is Π_1^1 -complete.

It remains to be shown that the property is in Π_1^1 (in [2] this has been done only reductions of length ω). A finite or infinite term t can be encoded as a function $t : \mathbb{N} \rightarrow \mathbb{N}$ (from positions to symbols from the signature). An infinite reduction can be rendered as a function $\sigma : \alpha \rightarrow ((\mathbb{N} \rightarrow \mathbb{N}) \times \mathbb{N})$ from an ordinal α to terms together with the rewrite position (here we assume that an ordinal is the set of all smaller ordinals) where $\sigma(\beta)$ is the β -th term of the sequence together with the rewrite position, and we require:

- (i) $\sigma(\beta)$ rewrites to $\sigma(\beta + 1)$ for all $\beta < \alpha$, and
- (ii) for all limit ordinals $\beta < \alpha$, and γ approaching β from below, we have:
 - $\sigma(\gamma)$ converges to $\sigma(\beta)$, and
 - the depth of the γ -th rewrite steps tends to infinity.

If condition (ii) holds for all limit ordinals $\beta \leq \alpha$ then the rewrite sequence σ is called strongly convergent. An ordinal α can be viewed as a well-founded relation $\alpha \subseteq \mathbb{N} \times \mathbb{N}$. The property of a relation to be well-founded can be expressed by a Π_1^1 -formula, and the above properties on rewrite sequences are arithmetic. By [4] the property $\text{SN}_R^\infty(s)$ holds if and only if all reductions admitted by s are strongly convergent. Hence SN_R^∞ and $\text{SN}_R^\infty(s)$ can be expressed by a Π_1^1 -formula since the above conditions (i) and (ii) are arithmetic. □

Using a minor modification of the term rewriting system from Proposition 6 we obtain that weak normalization for single terms and *reachability* are Σ_1^1 -complete, that is, the problem of deciding on the input of a TRS S and terms s, t whether $s \twoheadrightarrow_S t$.

Theorem 8. *Infinitary weak normalization for single terms, $\text{WN}_R^\infty(s)$, and reachability in infinitary rewriting are Σ_1^1 -complete.*

Proof. Let M be an arbitrary Turing machine. We define the TRS S' together to consist of the rules of $R_M^\bullet \uplus R_{\text{pickn}}$ together with:

$$\text{run}(T, \text{ok}(x), \text{ok}(y)) \rightarrow \bullet(\text{run}(q_0(x, y), \text{ok}(y), \text{pickn})), \quad (2)$$

and define a term $t := \text{run}(\mathbf{T}, \text{pickn}, \text{pickn})$. We have $t \twoheadrightarrow_{S'} \bullet^\infty$ if and only if t admits a rewrite sequence containing infinitely many root steps with respect to the rewrite system S from Proposition 6. As a consequence we have:

$$t \twoheadrightarrow_{S'} \bullet^\infty \iff \neg \text{SN}_S^\infty(t) \iff >_{\mathbf{M}} \text{ is not well-founded.}$$

Hence reachability is Σ_1^1 -hard.

We add one more rule to S' :

$$\text{run}(x, y, z) \rightarrow \text{run}(x, y, z), \quad (3)$$

Note that this rule has no impact on reachability. Then $\text{WN}_{S'}^\infty(t)$ holds if and only if $t \twoheadrightarrow_{S'} \bullet^\infty$, and hence $\text{WN}_{S'}^\infty(t)$ is Σ_1^1 -complete.

Moreover, weak normalization for single terms and reachability are in Σ_1^1 . We have $\text{WN}_{S'}^\infty(t)$ if and only if there exists a normal form t' such that $t \twoheadrightarrow t'$, and we have reachability $s \twoheadrightarrow t$ if and only if there exists a reduction from s to t . The quantification over terms and rewrite sequences are existential set or function quantifiers (which can be compressed to one single quantifier), and all other properties are arithmetic; see the encoding of reduction sequences see the proof of Theorem 7. $\square$

2 Uniform Infinitary Weak Normalization

For $\sigma \in \Gamma^\omega$ and $i \in \mathbb{N}$ we write $\sigma_{<i}$ for the prefix of σ up to (excluding) position i , and $\sigma_{\geq i}$ for the suffix of σ starting from (including) position i . We define non-deterministic Turing machines with one-sided infinite tape.

Definition 9. A *non-deterministic (one-sided) Turing machine* $\mathbf{M}$ is a quadruple $\langle Q, \Gamma, q_0, \delta \rangle$ consisting of:

- finite set of states Q ,
- an initial state $q_0 \in Q$,
- a finite alphabet Γ containing a designated symbol $\square$, called *blank*, and
- a partial *transition function* $\delta : Q \times \Gamma \rightarrow \mathcal{P}(Q \times \Gamma \times \{L, R\})$.

A *configuration* of $\mathbf{M}$ is a triple $\langle q, \sigma, i \rangle$ consisting of a state $q \in Q$, a tape content $\sigma : \Gamma^\omega$, and the position of the head $i \in \mathbb{N}$.

For two configurations we define $\langle q, \sigma, i \rangle \rightarrow_{\mathbf{M}} \langle q', \sigma', i' \rangle$ whenever:

- $\langle q', f, L \rangle \in \delta(q, \sigma(i))$, $i > 0$ $i' = i - 1$, and $\sigma' = \sigma_{<i} f \sigma_{\geq i+1}$
- $\langle q', f, R \rangle \in \delta(q, \sigma(i))$, $i' = i + 1$, and $\sigma' = \sigma_{<i} f \sigma_{\geq i+1}$

An infinite sequence of configurations $r : \{\langle q_j, \sigma_j, i_j \rangle\}_{j \geq 0}$ is a *run of $\mathbf{M}$ on σ* if:

- (i) $\langle q_0, \sigma_0, i_0 \rangle = \langle q_0, \sigma, 0 \rangle$, and
- (ii) $\langle q_j, \sigma_j, i_j \rangle \rightarrow_{\mathbf{M}} \langle q_{j+1}, \sigma_{j+1}, i_{j+1} \rangle$ for all $j \geq 0$.

A run $r : \{\langle q_j, \sigma_j, i_j \rangle\}_{j \geq 0}$ is called *complete* if every position is visited, that is, $\forall n \geq 0. \exists j \geq 0. i_j = n$, and r is called *oscillating* if $\exists n \geq 0. \forall j \geq 0. \exists j' > j. i_{j'} = n$.

Definition 10. A run is called *accepting* if it is complete and non-oscillating. The ω -language $\mathcal{L}^\omega(\mathbf{M})$ accepted by a non-deterministic Turing machine $\mathbf{M}$ is:

$$\mathcal{L}^\omega(\mathbf{M}) = \{w \in \Gamma^\omega \mid \text{there exists an accepting run of } \mathbf{M} \text{ on } w\}$$

Notice that accepting runs visits every symbol at least once, but only finitely often. The following is a proposition from [1]:

Proposition 11 ([1]). *The set $\{\mathbf{M} \mid \mathcal{L}^\omega(\mathbf{M}) = \Gamma^\omega\}$ is Π_2^1 -complete.* $\square$

We use the translation of Turing machines $\mathbf{M}$ to string rewriting systems $S_{\mathbf{M}}$ from [5].

Definition 12. For every (non-deterministic) Turing machine $\mathbf{M} = \langle Q, \Gamma, q_0, \delta \rangle$ we define a TRS $S_{\mathbf{M}}$ as follows. The signature Σ consists of symbols from $Q \cup \Gamma$ all having arity 1. The rewrite rules of $S_{\mathbf{M}}$ are:

$$\begin{aligned} q(f(x)) &\rightarrow f'(q'(x)) && \text{for every } \langle q', f', R \rangle \in \delta(q, f) \\ g(q(f(x),)) &\rightarrow q'(g(f'(x))) && \text{for every } \langle q', f', L \rangle \in \delta(q, f) \end{aligned}$$

Definition 13. Let $\mathbf{M}$ be a non-deterministic Turing machine. We define a mapping $\varphi : (\Gamma \cup Q)^* \rightarrow \text{Ter}^\infty(\Gamma \cup Q)$ by $\varphi(aw) := a(\varphi(t))$, and we extend φ to configurations $\langle q, \sigma, i \rangle$ of $\mathbf{M}$ by defining: $\varphi(\langle q, \sigma, i \rangle) = \varphi(\sigma_{<i} q \sigma_{\geq i})$.

The following lemma follows immediately from the definition of $S_{\mathbf{M}}$:

Lemma 14. *Let $\mathbf{M}$ be a non-deterministic Turing machine. For configurations c_1, c_2 of $\mathbf{M}$ we have $c_1 \rightarrow_{\mathbf{M}} c_2$ if and only if $\varphi(c_1) \rightarrow_{S_{\mathbf{M}}} \varphi(c_2)$.* $\square$

Then we obtain the following lemma establishing a correspondence of strongly convergent rewrite sequences and complete, non-oscillating runs:

Lemma 15. *Let $\mathbf{M}$ be a non-deterministic Turing machine and $\sigma \in \Gamma^\omega$. Then $\sigma \in \mathcal{L}^\omega(\mathbf{M})$ if and only if $q_0(\varphi(\sigma)) \twoheadrightarrow_{S_{\mathbf{M}}} t$ for some ground term $t \in \text{Ter}^\infty(\Gamma)$, that is, $t = \varphi(\sigma')$ for some $\sigma' \in \Gamma^\omega$.*

Proof. By Lemma 14 every rewrite sequence $\varphi(\langle q_0, \sigma, 0 \rangle) \twoheadrightarrow \dots$ corresponds to a run of $\mathbf{M}$ on σ . By definition of $\twoheadrightarrow$ the limit term exists if and only if the rewrite sequence is strongly convergent and this holds if and only if every rewrite position occurs at only finitely often, that is, the run is complete and non-oscillating. $\square$

Theorem 16. *Uniform infinitary weak normalization, WN_R^∞ , is Π_2^1 -complete.*

Proof. Let $\mathbf{M}$ be a non-deterministic Turing machine. We define the TRS R as an extension of the TRS $S_{\mathbf{M}}$ with the following rules:

$$\text{run}(x, y, z, z) \rightarrow \text{run}(\xi, q_0(z), \Delta_1(z), \Delta_1(z)) \quad (4)$$

$$\text{run}(x, x, y, z) \rightarrow \perp \quad (5)$$

$$q(x) \rightarrow \perp \quad \text{for } q \in Q \quad (6)$$

$$\xi \rightarrow f(\xi) \quad \text{for all } f \in \Gamma \quad (7)$$

$$\Delta_1(f(x)) \rightarrow f(\Delta_1(x)) \quad \text{for all } f \in \Gamma \quad (8)$$

$$\Delta_2(f(x)) \rightarrow f(\Delta_2(x)) \quad \text{for all } f \in \Gamma \quad (9)$$

The rules for ξ , Δ_1 and Δ_2 are obviously infinitary normalizing, in particular the normal forms of ξ are exactly all ground terms from $Ter^\infty(\Gamma)$. By application of $q(x) \rightarrow \perp$ every term can be rewritten to a normal form with respect to S_M . Moreover, $\Delta_1(t)$ and $\Delta_2(t)$ have a common reduct if and only if $t \twoheadrightarrow t'$ for a ground term from $Ter^\infty(\Gamma)$.

Assume that $\mathcal{L}^\omega(M) = \Gamma^\omega$. If there exists a term that is not infinitary weakly normalizing, then by the above considerations it must admit a rewrite sequence where (at some fixed position) the first rule is applied infinitely often. By the shape of (4) $run(s, t, u, u) \rightarrow run(\xi, q_0(u), \Delta_1(u), \Delta_1(u))$ this can only occur if $\Delta_1(u)$ and $\Delta_2(u)$ have a common reduct. This implies that $u \twoheadrightarrow u'$ for some ground term $u' \in Ter^\infty(\Gamma)$. Then $q_0(u) \twoheadrightarrow q_0(u')$ and by Lemma 15 we have $q_0(u') \twoheadrightarrow u''$ for some ground term $u'' \in Ter^\infty(\Gamma)$ (note that $\varphi^{-1}(u') \in \mathcal{L}^\omega(M)$). Since also $\xi \twoheadrightarrow u''$ we get a rewrite step of the form $run(u'', u'', \xi, u) \rightarrow \perp$. Hence every term is $WN_{S_M}^\infty$.

Assume that $\mathcal{L}^\omega(M) \neq \Gamma^\omega$. Then there exists $w \in \Gamma^\omega$ for which there exists no accepting run of M . Let $u = \varphi(w)$. We claim that the term

$$run(\xi, q_0(u), \Delta_1(u), \Delta_2(u))$$

is not infinitary weakly normalizing. Note that $q_0(u)$ does not reduce to a term containing ξ , and the only ξ -free reducts of ξ are terms from $Ter^\infty(\Gamma)$. However, $q_0(u)$ does not reduce to a ground terms from $Ter^\infty(\Gamma)$ by Lemma 15 since there M admits no accepting run for w . Consequently, the rule 5 is never applicable, and we cannot get rid of the redexes in $\Delta_1(u)$ and $\Delta_2(u)$ unless we reduce both to their unique normal form: $\Delta_1(u) \twoheadrightarrow u$ and $\Delta_2(u) \twoheadrightarrow u$. However, then we have a root redex giving rise to a step:

$$run(s, t, u, u) \rightarrow run(\xi, q_0(u), \Delta_1(u), \Delta_1(u))$$

This concludes Π_2^0 -hardness of uniform WN^∞ .

It remains be shown that WN^∞ is in Π_2^1 . The property WN_R^∞ holds if and only if for all terms $s \in Ter^\infty(\Sigma)$ there exists a reduction to a normal form. (For the encoding of reduction sequences see the proof of Theorem 7.) Hence WN^∞ is in Π_2^1 . $\square$

References

1. J. Castro and F. Cucker. Nondeterministic ω -computations and the analytical hierarchy. *Journal Math. Logik und Grundlagen d. Math.*, 35:333–342, 1989.
2. J. Endrullis, H. Geuvers, and H. Zantema. Degrees of undecidability in term rewriting. In E. Grädel and R. Kahle, editors, *Proceedings of Computer Science Logic (CSL09)*, volume 5771 of *Lecture Notes in Computer Science*, pages 255–270. Springer, 2009.
3. J. W. Klop. Term rewriting systems. In S. Abramsky, D. M. Gabbay, and S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, volume 2, pages 1–116. Oxford University Press, Inc., 1992.

4. J. W. Klop and R. C. d. Vrijer. Infinitary Normalization. In S. Artemov, H. Baringer, A. d'Avila Garcez, L. Lamb, and J. Woods, editors, *We Will Show Them: Essays in Honour of Dov Gabbay*, volume 2, pages 169–192. College Publ., 2005.
5. Terese. *Term Rewriting Systems*, volume 55 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2003.